

LA-UR-21-24865

Approved for public release; distribution is unlimited.

Title: Semiconductor Nanocrystals: Tiny Particles with “Quantum Powers”

Author(s): Klimov, Victor Ivanovich

Intended for: Materials Capability Review

Issued: 2021-05-19

Disclaimer:


Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



Semiconductor Nanocrystals: Tiny Particles with “Quantum Powers”

Victor I. Klimov,
Chemistry Division, C-PCS

Los Alamos Quantum Dots: Twenty Five Years of Innovation...and Fun



Materials for the Future

■ Center for Integrated Nanotechnologies (CINT), DOE's NSRC (2000 - ...)



■ Center for Advanced Solar Photophysics (CASP), DOE's EFRC (2009 - 2019)



Materials for the Future

■ LANL's Pillar: Materials for the future


Controlled functionality through:

Quantum confinement

Wavefunction engineering

*Engineered surfaces &
interfaces*

Controlled inter-dot coupling



■ LANL's Areas of Leadership

Integrated nanomaterials:

*Emergent functionality at the
nanoscale*

*Next generation materials for
manipulating photons*

*Advanced materials for
optoelectronics*


■ Overarching Science Questions:

How to predict & control design of nanostructure and/or interfaces to achieve emergent functionalities?

How do we understand & exploit competing interactions associated with reduced dimensionality & interfaces to yield tailored properties?




Born in Colloidal Solution: Artificial Atoms with Tunable Energies

■ Fabrication: Moderate-temperature colloidal chemistry



$$R = 10\text{--}50 \text{ \AA}, \Delta R/R = 4\text{--}7\%$$

■ Size-tunable electronic energies



$$E_g(NQD) = E_g(bulk) +$$

$$+ \frac{\hbar^2 \pi^2}{2m_{eh} R^2}$$


↑
quantum
confinement term
($\geq 200 \text{ meV}$)

Tiny Particles...with Magic Powers


■ Lighting & Displays




■ Lasing




■ Solar energy




■ Sensing



■ Electronics





LDRD DR Project: Quantum Photonics with Semiconductor Nanocrystals

■ **Overarching Objective:** To explore the utility of colloidal quantum dots as highly-versatile, solution-processible materials for implementing wavelength-selectable single-photon emitters (SPEs) complemented by on-chip light-amplification circuits, laser sources, and logic-gate devices.

■ **'Single-photon emitter' (SPE):** Emits exactly **one photon at a designated time** and all photons are quantum-mechanically identical or **'indistinguishable'**

Chemistry:

C-PCS

Materials Physics & Applications:

Nat High-Magnetic Field Lab


Center for Integrated Nanotech


Nuclear Eng & Nonproliferation:

NEN-1

Theory:

T-1








Project Milestones & Timeline




Project Timeline

- Y1 **“Quantum light 101”**
Achieving high single-photon purity, near-infrared (NIR) wavelength, high spectral stability
- Y2 **“Quantum light 201”**
Expanding/enhancing QL101 capabilities: electrical pumping, coupling to cavities...
- Y3 **“Quantum light 301”**
Achieving high excitonic coherence, photon indistinguishability, maintained at elevated temperature

Prospective applications: High single-photon purity




Prospective applications: High excitonic coherence




Photon Statistics & Single-Photon Purity

Photon statistics: Chaotic (classical), coherent (laser) & anti-bunched (quantum) light

Hanbury, Twiss & Brown experiment






Ideal single-photon emitter: $g^{(2)}(0) = 0$



Antibunching mechanisms in colloidal QDs: Quantized Auger recombination

$$F_{\text{spp}} = 1 - g^{(2)}(0)/g^{(2)}(T) = 1 - Q_{\text{XX}}/Q_{\text{X}} = 1 - 4\tau_{\text{XX}}/\tau_{\text{X}} = 0.99$$



V. Klimov *et al.*,
Science **287**, 1011
(2000)





Pros & Cons of “Giant” CdSe/CdS QDs

■ “Giant” thick-shell CdSe/CdS QDs: Nearly complete suppression of intensity fluctuations

C. Galand, et al.
Nature **479**, 203
(2011)





■ ... Still significant spectral fluctuations



Fixing the Problem of Standard “Giant” QDs


Asymmetrically strained, continuously graded CdSe/ZnCdSe cg-QDs



O. Kozlov... V.I. Klimov, *Science* 365, 672 (2019)

Sub-kT spectral fluctuations

J. Lim, Y-H. Park, V.I. Klimov,
Nature Mater. 18, 249 (2019)



CdSe/HgS/CdS QDs: Near-IR Upgrade for CdSe/CdS QDs

■ “Nonblinking” NIR CdSe/HgS/CdS QDs with atomically defined HgS interlayer


nature
nanotechnology

ARTICLES
<https://doi.org/10.1038/s41565-021-00871-x>
Check for updates

Highly versatile near-infrared emitters based on an atomically defined HgS interlayer embedded into a CdSe/CdS quantum dot




Vladimir Sayevich¹, Zachary L. Robinson^{1,2}, Younghee Kim¹, Oleg V. Kozlov¹, Heeyoung Jung¹, Tom Nakotte^{1,3}, Young-Shin Park^{1,4} and Victor I. Klimov^{1,3}

■ Thermodynamically controlled L-b-L growth





■ Atomic-level control of the HgS interlayer thickness

V. Sayevich, ... V.I. Klimov,
Nature Nanotech. March 25 (2021)




CdSe/HgS/CdS QDs: Perfect NIR Emitters

- Highly efficient, spectrally tunable NIR emitters (QY > 60%)




- Excellent electroluminescent materials (NIR-LEDs)



- Highly stable single-photon emitters


Nearly blinking-free emission

$$f_{ON} > 95\%$$




Perfect single-photon purity

$$F_{spp} > 97\%$$




Photon Amplification

- High-sensitivity detection using single-photon QD amplifiers (sub-threshold laser)




- cg-QDs are excellent optical gain materials (suppressed Auger decay, large gain)



O. Kozlov, ... V.I. Klimov,
Science **365**, 672 (2019)

J. Lim, Y.S. Park, V.I. Klimov,
Nature Mater. **17**, 42 (2018)

- Complete 1S and 1P population inversion using pulsed, current focusing QD-LEDs




H. Jung, Y.S. Park, V.I. Klimov,
Nature Phot. **in press** (2021)

Ongoing & Future Research

Primary Challenge: Achieve photon indistinguishability via control of extrinsic and intrinsic mechanisms for dephasing

Path forward

- Manipulation of NC electronic wavefunctions, surfaces, and matrices to eliminate spectral instabilities due to fluctuating environment
- Manipulation of NC electronic states, phonon spectra and matrices to suppress dephasing due to electron-phonon scattering
- Exploiting resonant coupling to optical cavities to enhance and sharpen the coherent exciton line and suppress incoherent phonon replicas.



Acknowledgements



■ Nanotechnology & Advanced Spectroscopy Team



■ Quantum Photonics LDRD-DR Project



Victor Klimov
(C-PCS)



Sergei Tretiak
(T-1)



Sergei Ivanov
(MPA-CINT)

C-PCS
MPA-MAG
MPA-CINT
NEN-1
T-1



Mark Croce
(NEN-1)



Scott Crooker
(MPA-MAG)



Robinson (PD)



Sayevich (PD)



Kim (PD)



Ghosh (PD)





Ahn (PD)



Colloidal Quantum Dots as Single-Photon Emitters

- InAs/GaAs QDs ($T = 4.5 \text{ K}$)
Single-photon purity = 99.1%
Indistinguishability = 97.5%
 Dusanowski, et al., *Phys. Rev. Lett.* **122**, 173602 (2019)



- Free-standing particles: uniquely suited for applications in imaging and sensing (by integrating with, e.g., scan probes)
- Easily scalable structures prepared using non-clean room techniques (e.g., inkjet printing)
- Very inexpensive

Epitaxial quantum dots

- Small inter-level spacing: Need cryogenic-cooling (liquid helium)
- Limited range of accessible wavelengths ($\sim 800 - 1100 \text{ nm}$; defined by InGaAs band-gap)
- Difficulty for on-chip integration with Si circuits (lattice mismatch)
- Dot-to-dot variability in emission energy (size polydispersity)

Colloidal quantum dots

- Can serve as SPEs at elevated T : large inter-level spacing ($> k_B T$), strongly confined acoustic phonons
- Readily tunable emission *via* size-controlled band-gap (UV – visible – IR)
- Chemically processable particles: easy to integrate with electronic and photonic circuits
- In principle, can be prepared as monodispersed structures with identical emission energies